

# Linear Algebra Review

(Adapted from Punit Shah's [slides](#))

Introduction to Machine Learning (CSC 2515)  
Fall 2022

University of Toronto

# Basics

- A scalar is a number.
- A vector is a 1-D array of numbers. The set of vectors of length  $n$  with real elements is denoted by  $\mathbb{R}^n$ .
  - Vectors can be multiplied by a scalar.
  - Vectors can be added together if dimensions match.
- A matrix is a 2-D array of numbers. The set of  $m \times n$  matrices with real elements is denoted by  $\mathbb{R}^{m \times n}$ .
  - Matrices can be added together or multiplied by a scalar.
  - We can multiply Matrices to a vector if dimensions match.
- In the rest we denote scalars with lowercase letters like  $a$ , vectors with bold lowercase  $\mathbf{v}$ , and matrices with bold uppercase  $\mathbf{A}$ .

- Norms measure how “large” a vector is. They can be defined for matrices too.
- The  $\ell_p$ -norm for a vector  $\mathbf{x}$ :


$$\|\mathbf{x}\|_p = \left[ \sum_i |x_i|^p \right]^{\frac{1}{p}}.$$

- The  $\ell_2$ -norm is known as the Euclidean norm.
- The  $\ell_1$ -norm is known as the Manhattan norm, i.e.,  $\|\mathbf{x}\|_1 = \sum_i |x_i|$ .
- The  $\ell_\infty$  is the max (or supremum) norm, i.e.,  $\|\mathbf{x}\|_\infty = \max_i |x_i|$ .

# Dot Product

- Dot product is defined as  $\mathbf{v} \cdot \mathbf{u} = \mathbf{v}^\top \mathbf{u} = \sum_i u_i v_i$ .
- The  $\ell_2$  norm can be written in terms of dot product:  $\|\mathbf{u}\|_2 = \sqrt{\mathbf{u} \cdot \mathbf{u}}$ .
- Dot product of two vectors can be written in terms of their  $\ell_2$  norms and the angle  $\theta$  between them:

$$\mathbf{a}^\top \mathbf{b} = \|\mathbf{a}\|_2 \|\mathbf{b}\|_2 \cos(\theta).$$



# Cosine Similarity

- Cosine between two vectors is a measure of their similarity:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|}.$$

- **Orthogonal Vectors:** Two vectors  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal to each other if  $\mathbf{a} \cdot \mathbf{b} = 0$ .

# Vector Projection

- Given two vectors  $\mathbf{a}$  and  $\mathbf{b}$ , let  $\hat{\mathbf{b}} = \frac{\mathbf{b}}{\|\mathbf{b}\|}$  be the unit vector in the direction of  $\mathbf{b}$ .
- Then  $\mathbf{a}_1 = a_1 \cdot \hat{\mathbf{b}}$  is the orthogonal projection of  $\mathbf{a}$  onto a straight line parallel to  $\mathbf{b}$ , where

$$a_1 = \|\mathbf{a}\| \cos(\theta) = \mathbf{a} \cdot \hat{\mathbf{b}} = \mathbf{a} \cdot \frac{\mathbf{b}}{\|\mathbf{b}\|}$$




Image taken from [wikipedia](#).

- Trace is the sum of all the diagonal elements of a matrix, i.e.,

$$\text{Tr}(\mathbf{A}) = \sum_i A_{i,i}.$$

- Cyclic property:

$$\text{Tr}(\mathbf{ABC}) = \text{Tr}(\mathbf{CAB}) = \text{Tr}(\mathbf{BCA}).$$


# Multiplication

- Matrix-vector multiplication is a linear transformation. In other words,

$$\mathbf{M}(v_1 + av_2) = \mathbf{M}v_1 + a\mathbf{M}v_2 \implies (\mathbf{M}v)_i = \sum_j M_{i,j}v_j.$$

- Matrix-matrix multiplication is the composition of linear transformations, i.e.,

$$(\mathbf{A}\mathbf{B})v = \mathbf{A}(\mathbf{B}v) \implies (\mathbf{A}\mathbf{B})_{i,j} = \sum_k A_{i,k}B_{k,j}.$$





# Invertibility

- $\mathbf{I}$  denotes the identity matrix which is a square matrix of zeros with ones along the diagonal. It has the property  $\mathbf{IA} = \mathbf{A}$  ( $\mathbf{BI} = \mathbf{B}$ ) and  $\mathbf{Iv} = \mathbf{v}$
- A square matrix  $\mathbf{A}$  is invertible if  $\mathbf{A}^{-1}$  exists such that  $\mathbf{A}^{-1}\mathbf{A} = \mathbf{AA}^{-1} = \mathbf{I}$ .
- Not all non-zero matrices are invertible, e.g., the following matrix is not invertible:

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

# Transposition

- Transposition is an operation on matrices (and vectors) that interchange rows with columns.  $(\mathbf{A}^\top)_{i,j} = \mathbf{A}_{j,i}$ .
- $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$ .
- $\mathbf{A}$  is called symmetric when  $\mathbf{A} = \mathbf{A}^\top$ .
- $\mathbf{A}$  is called orthogonal when  $\mathbf{AA}^\top = \mathbf{A}^\top \mathbf{A} = \mathbf{I}$  or  $\mathbf{A}^{-1} = \mathbf{A}^\top$ .

# Diagonal Matrix

- A diagonal matrix has all entries equal to zero except the diagonal entries which might or might not be zero, e.g. identity matrix.
- A square diagonal matrix with diagonal entries given by entries of vector  $\mathbf{v}$  is denoted by  $\text{diag}(\mathbf{v})$ .
- Multiplying vector  $\mathbf{x}$  by a diagonal matrix is efficient:

$$\text{diag}(\mathbf{v})\mathbf{x} = \mathbf{v} \odot \mathbf{x},$$

where  $\odot$  is the entrywise product.

- Inverting a square diagonal matrix is efficient

$$\text{diag}(\mathbf{v})^{-1} = \text{diag}\left(\left[\frac{1}{v_1}, \dots, \frac{1}{v_n}\right]^\top\right).$$

# Determinant

- Determinant of a square matrix is a mapping to scalars.

$$\det(\mathbf{A}) \quad \text{or} \quad |\mathbf{A}|$$

- Measures how much multiplication by the matrix expands or contracts the space.
- Determinant of product is the product of determinants:

$$\det(\mathbf{AB}) = \det(\mathbf{A})\det(\mathbf{B})$$

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

# List of Equivalencies

Assuming that  $\mathbf{A}$  is a square matrix, the following statements are equivalent

- $\mathbf{Ax} = \mathbf{b}$  has a **unique** solution (for every  $b$  with correct dimension).
- $\mathbf{Ax} = \mathbf{0}$  has a unique, trivial solution:  $\mathbf{x} = \mathbf{0}$ .
- Columns of  $\mathbf{A}$  are linearly independent.
- $\mathbf{A}$  is invertible, i.e.  $\mathbf{A}^{-1}$  exists.
- $\det(\mathbf{A}) \neq 0$

# Zero Determinant

If  $\det(\mathbf{A}) = 0$ , then:

- $\mathbf{A}$  is linearly dependent.
- $\mathbf{Ax} = \mathbf{b}$  has infinitely many solutions or no solution. These cases correspond to when  $\mathbf{b}$  is in the span of columns of  $\mathbf{A}$  or out of it.
- $\mathbf{Ax} = \mathbf{0}$  has a non-zero solution. (since every scalar multiple of one solution is a solution and there is a non-zero solution we get infinitely many solutions.)

# Matrix Decomposition

- We can decompose an integer into its prime factors, e.g.,  
 $12 = 2 \times 2 \times 3$ .
- Similarly, matrices can be decomposed into product of other matrices.

$$\mathbf{A} = \mathbf{V}\text{diag}(\boldsymbol{\lambda})\mathbf{V}^{-1}$$

- Examples are Eigendecomposition, SVD, Schur decomposition, LU decomposition, . . . .

# Eigenvectors

- An eigenvector of a square matrix  $\mathbf{A}$  is a nonzero vector  $\mathbf{v}$  such that multiplication by  $\mathbf{A}$  only changes the scale of  $\mathbf{v}$ .

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$$

- The scalar  $\lambda$  is known as the **eigenvalue**.
- If  $\mathbf{v}$  is an eigenvector of  $\mathbf{A}$ , so is any rescaled vector  $s\mathbf{v}$ . Moreover,  $s\mathbf{v}$  still has the same eigenvalue. Thus, we constrain the eigenvector to be of unit length:

$$\|\mathbf{v}\|_2 = 1$$



# Characteristic Polynomial(1)

- Eigenvalue equation of matrix  $\mathbf{A}$ .

$$\begin{aligned}\mathbf{A}\mathbf{v} &= \lambda\mathbf{v} \\ \lambda\mathbf{v} - \mathbf{A}\mathbf{v} &= \mathbf{0} \\ (\lambda\mathbf{I} - \mathbf{A})\mathbf{v} &= \mathbf{0}\end{aligned}$$

- If nonzero solution for  $\mathbf{v}$  exists, then it must be the case that:

$$\det(\lambda\mathbf{I} - \mathbf{A}) = 0$$

- Unpacking the determinant as a function of  $\lambda$ , we get:

$$P_A(\lambda) = \det(\lambda\mathbf{I} - \mathbf{A}) = 1 \times \lambda^n + c_{n-1} \times \lambda^{n-1} + \dots + c_0$$

- This is called the characteristic polynomial of  $\mathbf{A}$ .

## Characteristic Polynomial(2)

- If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are roots of the characteristic polynomial, they are eigenvalues of  $\mathbf{A}$  and we have  $P_A(\lambda) = \prod_{i=1}^n (\lambda - \lambda_i)$ .
- $c_{n-1} = -\sum_{i=1}^n \lambda_i = -tr(A)$ . This means that the sum of eigenvalues equals to the trace of the matrix.
- $c_0 = (-1)^n \prod_{i=1}^n \lambda_i = (-1)^n det(\mathbf{A})$ . The determinant is equal to the product of eigenvalues.
- Roots might be complex. If a root has multiplicity of  $r_j > 1$  (This is called the algebraic dimension of eigenvalue), then the geometric dimension of eigenspace for that eigenvalue might be less than  $r_j$  (or equal but never more). But for every eigenvalue, one eigenvector is guaranteed.

## Example

- Consider the matrix:

$$\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

- The characteristic polynomial is:

$$\det(\lambda \mathbf{I} - \mathbf{A}) = \det \begin{bmatrix} \lambda - 2 & -1 \\ -1 & \lambda - 2 \end{bmatrix} = 3 - 4\lambda + \lambda^2 = 0$$

- It has roots  $\lambda = 1$  and  $\lambda = 3$  which are the two eigenvalues of  $\mathbf{A}$ .
- We can then solve for eigenvectors using  $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ :

$$\mathbf{v}_{\lambda=1} = [1, -1]^\top \quad \text{and} \quad \mathbf{v}_{\lambda=3} = [1, 1]^\top$$

# Eigendecomposition

- Suppose that  $n \times n$  matrix  $\mathbf{A}$  has  $n$  linearly independent eigenvectors  $\{\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(n)}\}$  with eigenvalues  $\{\lambda_1, \dots, \lambda_n\}$ .
- Concatenate eigenvectors (as columns) to form matrix  $\mathbf{V}$ .
- Concatenate eigenvalues to form vector  $\boldsymbol{\lambda} = [\lambda_1, \dots, \lambda_n]^\top$ .
- The **eigendecomposition** of  $\mathbf{A}$  is given by:

$$\mathbf{A}\mathbf{V} = \mathbf{V}\text{diag}(\boldsymbol{\lambda}) \implies \mathbf{A} = \mathbf{V}\text{diag}(\boldsymbol{\lambda})\mathbf{V}^{-1}$$


# Symmetric Matrices

- Every symmetric (hermitian) matrix of dimension  $n$  has a set of (not necessarily unique)  $n$  orthogonal eigenvectors. Furthermore, all eigenvalues are real.
- Every real symmetric matrix  $\mathbf{A}$  can be decomposed into real-valued eigenvectors and eigenvalues:

$$\mathbf{A} = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}^\top$$


- $\mathbf{Q}$  is an orthogonal matrix of the eigenvectors of  $\mathbf{A}$ , and  $\mathbf{\Lambda}$  is a diagonal matrix of eigenvalues.
- We can think of  $\mathbf{A}$  as scaling space by  $\lambda_i$  in direction  $\mathbf{v}^{(i)}$ .

Plot of unit vectors  $\mathbf{u} \in \mathbb{R}^2$   
(circle)



with two variables  $x_1$  and  $x_2$

Plot of vectors  $\mathbf{A}\mathbf{u}$   
(ellipse)



# Eigendecomposition is not Unique

- Decomposition is not unique when two eigenvalues are the same.
- By convention, order entries of  $\mathbf{\Lambda}$  in descending order. Then, eigendecomposition is unique if all eigenvalues have multiplicity equal to one.
- If any eigenvalue is zero, then the matrix is **singular**. Because if  $\mathbf{v}$  is the corresponding eigenvector we have:  $\mathbf{A}\mathbf{v} = 0\mathbf{v} = 0$ .

# Positive Definite Matrix

- If a symmetric matrix  $A$  has the property:

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} > 0 \quad \text{for any nonzero vector } \mathbf{x}$$

Then  $A$  is called **positive definite**.

- If the above inequality is not strict then  $A$  is called **positive semidefinite**.
- For positive (semi)definite matrices all eigenvalues are positive (non negative).

# Singular Value Decomposition (SVD)

- If  $\mathbf{A}$  is not square, eigendecomposition is undefined.
- **SVD** is a decomposition of the form  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$ .
- SVD is more general than eigendecomposition.
- Every real matrix has a SVD.



## SVD Definition (1)

- Write  $\mathbf{A}$  as a product of three matrices:  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$ .
- If  $\mathbf{A}$  is  $m \times n$ , then  $\mathbf{U}$  is  $m \times m$ ,  $\mathbf{D}$  is  $m \times n$ , and  $\mathbf{V}$  is  $n \times n$ .
- $\mathbf{U}$  and  $\mathbf{V}$  are orthogonal matrices, and  $\mathbf{D}$  is a diagonal matrix (not necessarily square).
- Diagonal entries of  $\mathbf{D}$  are called **singular values** of  $\mathbf{A}$ .
- Columns of  $\mathbf{U}$  are the **left singular vectors**, and columns of  $\mathbf{V}$  are the **right singular vectors**.

## SVD Definition (2)

- SVD can be interpreted in terms of eigendecomposition.
- Left singular vectors of  $\mathbf{A}$  are the eigenvectors of  $\mathbf{A}\mathbf{A}^\top$ .
- Right singular vectors of  $\mathbf{A}$  are the eigenvectors of  $\mathbf{A}^\top\mathbf{A}$ .
- Nonzero singular values of  $\mathbf{A}$  are square roots of eigenvalues of  $\mathbf{A}^\top\mathbf{A}$  and  $\mathbf{A}\mathbf{A}^\top$ .
- Numbers on the diagonal of  $D$  are sorted largest to smallest and are non-negative ( $\mathbf{A}^\top\mathbf{A}$  and  $\mathbf{A}\mathbf{A}^\top$  are semipositive definite.).

# Matrix norms

- We may define norms for matrices too. We can either treat a matrix as a vector, and define a norm based on an entrywise norm (example: Frobenius norm). Or we may use a vector norm to “induce” a norm on matrices.
- Frobenius norm:

$$\|A\|_F = \sqrt{\sum_{i,j} a_{i,j}^2}.$$

- Vector-induced (or operator, or spectral) norm:

$$\|A\|_2 = \sup_{\|x\|_2=1} \|Ax\|_2.$$

# SVD Optimality

- Given a matrix  $\mathbf{A}$ , SVD allows us to find its “best” (to be defined) rank- $r$  approximation  $\mathbf{A}_r$ .
- We can write  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$  as  $\mathbf{A} = \sum_{i=1}^n d_i \mathbf{u}_i \mathbf{v}_i^\top$ .
- For  $r \leq n$ , construct  $\mathbf{A}_r = \sum_{i=1}^r d_i \mathbf{u}_i \mathbf{v}_i^\top$ .
- The matrix  $\mathbf{A}_r$  is a rank- $r$  approximation of  $A$ . Moreover, it is the best approximation of rank  $r$  by many norms:
  - When considering the operator (or spectral) norm, it is optimal. This means that  $\|A - A_r\|_2 \leq \|A - B\|_2$  for any rank  $r$  matrix  $B$ .
  - When considering Frobenius norm, it is optimal. This means that  $\|A - A_r\|_F \leq \|A - B\|_F$  for any rank  $r$  matrix  $B$ . One way to interpret this inequality is that rows (or columns) of  $A_r$  are the projection of rows (or columns) of  $A$  on the best  $r$  dimensional subspace, in the sense that this projection minimizes the sum of squared distances.